

ZOETIS DRY COW TUBES STAND UP TO THE COMPETITION.

BRAND	SPECTRAMAST® DC*	ALBADRY PLUS® (penicillin G procaine and novobiocin sodium) Suspension	TOMORROW® Cefa-Dri® (cephapirin benzathine)	Orbenin®-DC (benzathine cloxacillin)	Dry-Clox® (benzathine cloxacillin)
ACTIVE INGREDIENT	Ceftiofur 500 mg	Penicillin 200,000 IU and novobiocin 400 mg	Cephapirin 300 mg	Cloxacillin 500 mg	Cloxacillin 500 mg
INDICATIONS	Treatment	Treatment of subclinical mastitis	Treatment of mastitis	Treatment and prophylaxis of mastitis	Treatment of mastitis
LABELED PATHOGENS	Staph. aureus Strep. dysgalactiae Strep. uberis	Staph. aureus Strep. agalactiae	Staph. aureus Strep. agalactiae	Staph. aureus Strep. agalactiae	Staph. aureus Strep. agalactiae
PRE-SLAUGHTER WITHDRAWAL*	16 days	30 days	42 days	28 days	30 days
MILK DISCARD**	0 hours	72 hours	72 hours	0 hours	0 hours
DRY PERIOD LENGTH	30 days	30 days	30 days	28 days	30 days
AVAILABILITY	Rx	OTC	OTC	Rx	Rx
YEAR INTRODUCED	2005	1983	1978	1975	1975

*After last administration (or treatment)

**Milk discard times begin at first milking post freshening and require completion of a minimum dry cow period.



KEY FEATURES:

- Attacks more major mastitis-causing pathogens, including *Staphylococcus aureus*, *Streptococcus dysgalactiae* and *Strep. uberis*
- Shortest meat withdrawal — allows you to maximize your management options
- Zero milk discard*** — so you can get them back in the milking string faster
- Greater flexibility in milk and cattle management decisions

Important Safety Information: Inappropriate dosage or treatment intervals with SPECTRAMAST DC and/or failure to complete a minimum dry cow period (30 days) may result in violative milk residues. In cows completing a 30-day dry cow period, no milk discard is necessary. Following treatment with SPECTRAMAST DC, a 16-day pre-slaughter withdrawal is required. As with all drugs, SPECTRAMAST DC should not be used in animals found to be hypersensitive to the product.

***Zero milk discard period after calving following a 30-day dry cow period.



KEY FEATURES:

- Unique combination of penicillin and novobiocin provides reliable therapy for subclinical mastitis in dry cows
- Has a synergistic effect on bacterial isolates from bovine intramammary infections¹
- Bactericidal activity against the two common mastitis-causing pathogens — *Staph. aureus* and *Strep. agalactiae*
- Helps eliminate single-antibiotic failures and reduces the likelihood of resistance

Important Safety Information: Do not use ALBADRY PLUS less than 30 days prior to calving. Milk from treated cows must not be used for food during the first 72 hours after calving. Treated animals must not be slaughtered for food for 30 days following udder infusion. See full Prescribing Information attached.

¹Wheeler SJ, Edmondson PW, et al. Effect of Penicillin/Novobiocin (TETRADELTA™ Dry Cow, ALBADRY PLUS® Sterile Suspension) Dry Cow Therapy on Somatic Cell Count of Dairy Cows Over the Dry Period. Proc International Buiatrics Congress 2000. All trademarks are the property of Zoetis Inc., its affiliates and/or its licensors. ©2013 Zoetis Inc. All rights reserved. GDR13164

SPECTRAMAST® DC


brand of cefotiofur hydrochloride sterile suspension

For Intramammary Infusion in Dry Dairy Cattle Only

FOR USE IN ANIMALS ONLY — NOT FOR HUMAN USE

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: Cefotiofur hydrochloride is a cephalosporin antibiotic.



Chemical Name of Cefotiofur Hydrochloride U-64279A *HCl

5-Thi-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[2-(2-amino-4-thiazoyl)-2-(methoxyimino)acylamino]-3-{(2-furylcarbonyl)thiomethyl}-8-oxo-hydrochloride.

Cefotiofur Hydrochloride Sterile Suspension is an oil-based sterile suspension.

Each 10 mL PLASTET® Disposable Syringe Contains:
Cefotiofur Equivalents (as the hydrochloride salt) 500 mg
Microcrystalline Wax 500 mg
Colloisens 4 g.

Table 1. Cefotiofur MIC values* for isolates from a multi-site clinical field study evaluating subclinical mastitis in dry dairy cows in the U.S. during 2000

Table 2. Cefotiofur MIC values* for mastitis pathogens from diagnostic laboratories in the U.S. and Canada

Organism No. Isolated Date MIC_{C₅₀}* (µg/mL) MIC range (µg/mL)

Staphylococcus aureus 300 1.0 ≤0.06

Streptococcus dysgalactiae 55 1.0 ≤0.06

Streptococcus uberis 58 1.0 ≤0.06

Cohlglass (-) 139 2000-2001 1.0 ≤0.06

Staphylococcus 15 1991-1992 1.0 0.13-2.0

Staphylococcus 10 1993 1.0 0.25-1.0

aureus 107 1995 1.0 0.25-1.0

Streptococcus 61 2000 1.0 ≤0.06

dysgalactiae 139 2000-2001 1.0 ≤0.06

Streptococcus 15 1991-1992 1.0 0.06-1.0

dysgalactiae 152 1997-1999 0.25 No range†

Streptococcus 64 2000 0.06 ≤0.06

uberis 22 1991-1992 0.5 ≤0.06

Streptococcus 133 1997-1999 0.5 ≤0.06

uberis 20 2000 1.0 ≤0.06

Escherichia coli 39 1991-1992 1.0 0.25-1.0

Escherichia coli 52 2000 0.5 ≤0.06

Escherichia coli 40 1993 0.5 0.13-1.0

Escherichia coli 52 2000 0.5 ≤0.06

* The above *in vitro* data are available, but their clinical significance is unknown.

** The MIC for 90% of the isolates.

† No range of isolates yielded the same value.

Based on pharmacokinetic, milk residue and clinical effectiveness studies in dairy cattle following intramammary infusion of cefotiofur and the MIC and disk (30 µg diffusion) dilution data from mastitis pathogens are recommended by the Clinical and Laboratories Standards Institute (CLSI) (Table 3).

Table 3. Current recommended interpretive criteria established by CLSI for cefotiofur for Bovine Mastitis

Bovine Mastitis Organisms Disk Content

Zone diameter (mm)

MIC breakpoint (µg/mL)

S I R S I R

Staphylococcus aureus

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Streptococcus dysgalactiae

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Streptococcus uberis

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Streptococcus agalactiae

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Escherichia coli

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

S - Susceptible I - Intermediate R - Resistant

Standardized procedures require the use of laboratory control organisms for both standardized diffusion techniques and standardized disk diffusion techniques. The 30 µg cefotiofur sodium reference powder (or disk) should provide the following MIC values for this reference strain. The cefotiofur sodium disks or standard reference powder is appropriate for cefotiofur hydrochloride (Table 4).

Table 4. Acceptable quality control ranges for cefotiofur against CLSI recommended American Type Culture Collection (ATCC) reference strains

Organism (ATCC No.) Zone diameter (mm)

MIC range (µg/mL)

Escherichia coli (25922) 26 to 31 0.25 to 1.0

Staphylococcus aureus (2923) 27 to 31 —

Pseudomonas aeruginosa (2783) 14 to 18 16.0 to 64.0

*All testing performed using a 30 µg disk.

†No information or to obtain a material safety data sheet.

‡Milk taken from cows completing a 30-day dry cow period may be used for food with no milk discard due to cefotiofur residues.

§No pre-slaughter withdrawal period is required for neonatal calves born from treated cows regardless of colostrum consumption.

3. Following intramammary infusion, a 16-day pre-slaughter withdrawal period is required for treated cows.

4. Use of this product in a manner other than indicated under DOSAGE might result in violative residues.

CLINICAL MICROBIOLOGY

Cefotiofur is a broad-spectrum cephalosporin antibiotic that exerts its effect by inhibiting bacterial cell wall synthesis. Like other beta-lactam antibiotics, the cephalosporins inhibit cell wall synthesis by interfering with the enzymes essential to peptidoglycan synthesis. This effect results in lysis of the bacterial cell and accounts for the bactericidal nature of these agents. Cefotiofur demonstrated *in vitro* activity against clinical isolates and isolates from diagnostic laboratories. The results of susceptibility testing of these isolates against cefotiofur are presented in Tables 1 and 2. Appropriate reference strains were also susceptible tested and their minimum inhibitory concentration (MIC) values and zone of inhibition with a 30 µg disk are presented in Table 4.

Table 1. Cefotiofur MIC values for isolates from a multi-site clinical field study evaluating subclinical mastitis in dry dairy cows in the U.S. during 2000

Table 2. Cefotiofur MIC values* for mastitis pathogens from diagnostic laboratories in the U.S. and Canada

Organism No. Isolated Date MIC_{C₅₀}* (µg/mL) MIC range (µg/mL)

Staphylococcus aureus 300 1.0 ≤0.06

Streptococcus dysgalactiae 55 1.0 ≤0.06

Streptococcus uberis 58 1.0 ≤0.06

Cohlglass (-) 139 2000-2001 1.0 ≤0.06

Staphylococcus 15 1991-1992 1.0 0.13-2.0

Staphylococcus 10 1993 1.0 0.25-1.0

aureus 107 1995 1.0 0.25-1.0

Streptococcus 61 2000 1.0 ≤0.06

dysgalactiae 139 2000-2001 1.0 ≤0.06

Streptococcus 15 1991-1992 1.0 0.06-1.0

dysgalactiae 152 1997-1999 0.25 No range†

Streptococcus 64 2000 0.06 ≤0.06

uberis 22 1991-1992 0.5 ≤0.06

Streptococcus 133 1997-1999 0.5 ≤0.06

uberis 20 2000 1.0 ≤0.06

Escherichia coli 39 1991-1992 1.0 0.25-1.0

Escherichia coli 52 2000 0.5 ≤0.06

Escherichia coli 40 1993 0.5 0.13-1.0

Escherichia coli 52 2000 0.5 ≤0.06

* The above *in vitro* data are available, but their clinical significance is unknown.

** The MIC for 90% of the isolates.

† No range of isolates yielded the same value.

Based on pharmacokinetic, milk residue and clinical effectiveness studies in dairy cattle following intramammary infusion of cefotiofur and the MIC and disk (30 µg diffusion) dilution data from mastitis pathogens are recommended by the Clinical and Laboratories Standards Institute (CLSI) (Table 3).

Table 3. Current recommended interpretive criteria established by CLSI for cefotiofur for Bovine Mastitis

Bovine Mastitis Organisms Disk Content

Zone diameter (mm)

MIC breakpoint (µg/mL)

S I R S I R

Staphylococcus aureus

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Streptococcus dysgalactiae

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Streptococcus uberis

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Streptococcus agalactiae

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

Escherichia coli

30 µg ≥21 18-20 ≤17 ≤2.0 4.0 ≥8.0

S - Susceptible I - Intermediate R - Resistant

Standardized procedures require the use of laboratory control organisms for both standardized diffusion techniques and standardized disk diffusion techniques. The 30 µg cefotiofur sodium reference powder (or disk) should give the following zone diameters and the cefotiofur sodium reference powder (or disk) should provide the following MIC values for this reference strain. The cefotiofur sodium disks or standard reference powder is appropriate for cefotiofur hydrochloride (Table 4).

Table 4. Acceptable quality control ranges for cefotiofur against CLSI recommended American Type Culture Collection (ATCC) reference strains

Organism (ATCC No.) Zone diameter (mm)

MIC range (µg/mL)

Escherichia coli (25922) 26 to 31 0.25 to 1.0

Staphylococcus aureus (2923) 27 to 31 —

Pseudomonas aeruginosa (2783) 14 to 18 16.0 to 64.0

*All testing performed using a 30 µg disk.

†No information or to obtain a material safety data sheet.

‡Milk taken from cows completing a 30-day dry cow period may be used for food with no milk discard due to cefotiofur residues.

§No pre-slaughter withdrawal period is required for neonatal calves born from treated cows regardless of colostrum consumption.

ALBADRY PLUS® Suspension

NDC 0009-3139-02

brand of penicillin G procaine and novobiocin sodium suspension

For the Treatment of Subclinical Mastitis in Dry Cows

For Udder Instillation in Dry Cows Only

FOR USE IN ANIMALS ONLY — NOT FOR HUMAN USE

Restricted Drug — Use Only as Directed (California)

DESCRIPTION

Each 10 mL PLASTET® Disposable Sy